



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/85, 5/16, C07H 15/12	A1	(11) International Publication Number: WO 92/15694 (43) International Publication Date: 17 September 1992 (17.09.92)
(21) International Application Number: PCT/US92/01899 (22) International Filing Date: 6 March 1992 (06.03.92) (30) Priority data: 666,252 8 March 1991 (08.03.91) US (71) Applicant: THE SALK INSTITUTE FOR BIOLOGICAL STUDIES [US/US]; 10010 North Torrey Pines Road, La Jolla, CA 92037 (US). (72) Inventors: WAHL, Geoffrey, M. ; 4258 Arguello Street, San Diego, CA 92103 (US). O'GORMAN, Stephen, V. ; 2761 Havasupai Avenue, San Diego, CA 92117 (US).		(74) Agents: CAMPBELL, Cathryn et al.; Pretty, Schroeder, Brueggemann & Clark, 444 South Flower Street, Suite 2000, Los Angeles, CA 90071 (US). (81) Designated States: AT (European patent), BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), MC (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>
(54) Title: FLP-MEDIATED GENE MODIFICATION IN MAMMALIAN CELLS, AND COMPOSITIONS AND CELLS USEFUL THEREFOR (57) Abstract A gene activation/inactivation and site-specific integration system has been developed for mammalian cells. The invention system is based on the recombination of transfected sequences by FLP, a recombinase derived from <i>Saccharomyces</i> . In several cell lines, FLP has been shown to rapidly and precisely recombine copies of its specific target sequence. For example, a chromosomally integrated, silent β -galactosidase reporter gene was activated for expression by FLP-mediated removal of intervening sequences to generate clones of marked cells. Alternatively, the reverse reaction can be used to target transfected DNA to specific chromosomal sites. These results demonstrate that FLP can be used, for example, to mosaically activate or inactivate transgenes for a variety of therapeutic purposes, as well as for analysis of vertebrate development.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCI on the front pages of pamphlets publishing international applications under the PCI.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

FLP-MEDIATED GENE MODIFICATION IN MAMMALIAN CELLS,
AND COMPOSITIONS AND CELLS USEFUL THEREFOR

This invention relates to recombinant DNA technology. In a particular aspect, this invention relates to methods for the site-specific recombination of DNA in mammalian cells or host mammalian organisms. In
5 another aspect, the present invention relates to novel DNA constructs, as well as compositions, cells and host organisms containing such constructs. In yet another aspect, the present invention relates to methods for the activation and/or inactivation of expression of
10 functional genes. In a further aspect, the present invention relates to methods for the introduction of DNA into specific sites in the genome of mammalian cells. In a still further aspect, the present invention relates to gene therapy methods. In still another aspect, the
15 present invention relates to means for the recovery of transfected DNA from a cell or host organism. In a still further aspect, the present invention relates to assay methods.

20

BACKGROUND OF THE INVENTION

Many recent manipulations of gene expression involve the introduction of transfected genes
25 (transgenes) to confer some novel property upon, or to alter some intrinsic property of, mammalian cells or organisms. The efficacy of such manipulations is often impaired by such problems as the inability to control the chromosomal site of transgene integration; or the
30 inability to control the number of copies of a transgene that integrate at the desired chromosomal site; or by difficulties in controlling the level, temporal characteristics, or tissue distribution of transgene

-2-

expression; or by the difficulty of modifying the structure of transgenes once they are integrated into mammalian chromosomes.

Transgenes are often introduced into mammalian cells or organisms to determine which components of a transgene are required for specific qualitative or quantitative alterations of the host system. Since both chromosomal position and copy number are major determinants of transgene function, the usefulness of these analyses is limited because current techniques for efficiently introducing transgenes into mammalian hosts result in the insertion of a variable number of transgene copies at random chromosomal positions. It is, therefore, difficult (if not impossible) to compare the effects of one transgene to those of another if the two transgenes occupy different chromosomal positions and are present in the genome at different copy numbers. Considerably more refined analyses would be possible if one could routinely introduce single copies of a variety of transgenes into a defined chromosomal position.

The spatial or temporal characteristics of transgene expression is difficult to control in intact organisms. The restricted expression of transgenes is potentially of great interest, as this technique can be employed for a variety of therapeutic applications, e.g., for the selective interruption of a defective gene, for the alteration of expression of a gene which is otherwise over-expressed or under-expressed, for the selective introduction of a gene whose product is desirable in the host, for the selective removal or disruption of a gene whose expression is no longer desired in the host, and the like.

Transgene expression is typically governed by a single set of control sequences, including promoters and enhancers which are physically linked to the

-3-

transgenes (i.e., cis-acting sequences). Considerably greater expression control could be achieved if transgene expression could be placed under the binary control of these cis-acting sequences, plus an additional set of sequences that were not physically linked to the transgenes (i.e., trans-acting sequences). A further advantage would be realized if the transient activity of these trans-acting functions resulted in a stable alteration in transgene expression. In this manner, it would be possible, for example, to introduce into a host a transgene whose expression would have lethal or deleterious effects if it was constitutively expressed in all cells. This would be accomplished by delaying the expression of the transgene to a specific time or developmental stage of interest, or by restricting the expression of the transgene to a specific subset of the cell population.

It is currently difficult (if not impossible) to precisely modify the structure of transgenes once they have been introduced into mammalian cells. In many applications of transgene technology, it would be desirable to introduce the transgene in one form, and to then be able to modify the transgene in a defined manner. By this means, transgenes could be activated or inactivated or the sequences which control transgene expression could be altered by either removing sequences present in the original transgene or by inserting additional sequences into the transgene.

Previous descriptions of recombinase-mediated rearrangement of chromosomal sequences in *Drosophila* and mammalian cells have not directly addressed the question of whether site-specific recombinases could routinely create a functional translational reading frame. Moreover, the reported efficiency of the prior art recombinase system, in the only other description of site-specific recombination in

-4-

mammalian cells reported to date [based on Cre recombinase, described by Sauer and Henderson in Nucleic Acids Research, Vol. 17: 147 (1989)] appears to be quite low.

5

BRIEF DESCRIPTION OF THE INVENTION

10 In accordance with the present invention, we have developed a system for the selective modification of chromosomal or extrachromosomal DNA in mammalian cells. Selective modification can involve the insertion of one DNA into another DNA (e.g., to create a hybrid gene, to activate a gene, to inactivate a gene, and the
15 like), or the removal of specific DNA molecule(s) from other DNA molecule(s) containing the DNA to be removed (e.g., to inactivate a gene, to bring desired DNA fragments into association with one another, and the like).

20 The recombination system of the present invention is based on site-specific recombinase, FLP. In one application of the invention recombination system, FLP-mediated removal of intervening sequences is required for the formation of a functional gene. Expression of
25 the functional gene therefore, falls under the control of both the regulatory sequences associated with the functional gene and also under the control of those sequences which direct FLP expression.

30 The reverse of the above-described process, i.e., the FLP-mediated introduction of DNA, provides a convenient and selective means to introduce DNA into specific sites in mammalian chromosomes.

35 FLP-mediated recombination of marker genes provides a means to follow the fate of various sequences over the course of development and/or from generation-to-generation. The recombination event creates a functional

-5-

marker gene. This gain-of-function system can be used for lineage analyses in a wide variety of tissues in different organisms. Prior to FLP-mediated recombination, the marker gene is normally silent, i.e., the phenotype typical of the marker is not observed. In the absence of FLP, spontaneous recombination to produce functional marker occurs only at very low frequencies. In the presence of FLP, functional marker is efficiently produced. In addition, this gain-of-function system is heritable and is easily detected by simple histochemical assays. For example, in transgenic mice, the lineages in which recombination is to occur can be controlled by appropriate selection of the promoters used to drive FLP expression. This could include promoters that are only transiently active at a developmental stage that substantially precedes overt cell differentiation. Since transcription of the marker gene is controlled by regulatory sequences associated therewith, functional marker genes can be expressed at later developmental stages, after cell differentiation has occurred. By this means, it is possible to construct a map for mammalian development that correlates embryonic patterns of gene expression with the organization of mature tissues.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 presents schematic diagrams of FLP-mediated recombination events. In Figure 1A, FLP-mediated introduction of DNA is illustrated, while in Figure 1B, FLP-mediated removal of intervening sequences is illustrated.

Figure 2 is presented in three parts. Figure 2A presents schematic diagrams of the expression vectors pFRT β GAL, pNEO β GAL, and pOG44 FLP. Figure 2B presents a Southern blot of Hirt lysates prepared from

-6-

293 (human embryonic kidney) cells transfected with one microgram of pNEO β GAL and varying amounts of the pOG44 FLP expression vector. Figure 2C graphically presents the β -galactosidase activities in the same transfections shown in part B, referred to above.

Figure 3A, at the top, presents a schematic of the pattern of plasmid integration in E25 deduced from Southern blot analysis. Figure 3A, in the middle, presents the predicted pattern for β -galactosidase positive subclones of E25 if precise recombination across the FLP-recombination target sites occurs. Figure 3A, at the bottom, presents the predicted pattern for β -galactosidase negative, neomycin resistant subclones of E25B2 after FLP mediated insertion of pOG45. Figure 3B presents an analysis of genomic DNA from a cell line with a single integrated copy of pNEO β GAL (i.e., CVNEO β GAL/E25, designated as E25), two derivative β -galactosidase-positive subclones (designated as E25B1 and E25B2), and two subclones derived from E25B2 after transfection with pOG45 (designated as B2N1 and B2N2).

DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention, there is provided a mammalian recombination system comprising:

- (i) FLP recombinase, or a nucleotide sequence encoding same, and
- (ii) a first DNA comprising a nucleotide sequence containing at least one FLP recombination target site.

In accordance with another embodiment of the present invention, there are provided novel DNA constructs useful for the introduction of DNA into the

-7-

genome of a transfected organism, said DNA construct comprising, as an autonomous fragment:

- (a) at least one FLP recombination target site,
- 5 (b) at least one restriction endonuclease recognition site,
- (c) at least one marker gene,
- (d) a bacterial origin of replication, and optionally
- 10 (e) a mammalian cellular or viral origin of DNA replication.

In accordance with yet another embodiment of the present invention, there are provided novel DNA constructs useful for the rescue of DNA from the genome of a transfected organism, said DNA construct comprising, 15 as an autonomous fragment, in the following order, reading from 5' to 3' along said fragment:

- (a) a first FLP recombination target site,
- (b) an insert portion comprising, in any 20 suitable sequence:
 - (1) at least one restriction endonuclease recognition site,
 - (2) at least one marker gene,
 - (3) a bacterial origin of replication, 25 and optionally
 - (4) a mammalian cellular or viral origin of DNA replication, and
- (c) a second FLP recombination target site in tandem with said first FLP recombination 30 target site.

In addition, there are provided methods for the recovery of transfected DNA from the genome of a transfected organism employing the above-described constructs.

In accordance with still another 35 embodiment of the present invention, there is provided a method for the assembly of a functional gene (which is

-8-

then suitable for activation of expression), in mammalian cells, by recombination of individually inactive gene segments derived from one or more gene(s) of interest, wherein each of said segments contains at least one recombination target site, said method comprising:

- 5 contacting said individually inactive gene segments with a FLP recombinase, under conditions suitable for recombination to occur, thereby providing a DNA sequence which encodes a functional gene of interest.

10 In accordance with a further embodiment of the present invention, there is provided a method for the disruption of functional gene(s) of interest, thereby inactivating expression of such gene(s), in mammalian cells, wherein said gene(s) of interest contain at least one FLP recombination target site, said method comprising contacting said gene(s) of interest with:

- 15 (i) a DNA segment which contains at least one FLP recombination target site, and
- 20 (ii) FLP recombinase;
- wherein said contacting is carried out under conditions suitable for recombination to occur between said gene and said DNA segment, thereby disrupting the gene(s) of interest and rendering said gene(s) non-functional.

25 In accordance with a still further embodiment of the present invention, there is provided a method for the precisely targeted integration of DNA into the genome of a host organism, said method comprising:

- 30 (i) introducing a FLP recombination target site into the genome of cells which are compatible with the cells of the subject,
- 35 (ii) introducing a first DNA comprising a nucleotide sequence containing at least one FLP recombination target site therein into the FLP recombination target site in the genome of said cells by contacting

-9-

said cells with said first DNA and FLP recombinase, and thereafter

(iii) introducing the cells produced by the process of step (ii) into said subject, wherein the resulting cells and/or organism have the optional ability to have DNA reproducibly and repetitively inserted into and/or recovered from the host cells and/or organism.

10 In accordance with another aspect of the present invention, there are provided mammalian cells, wherein the genomic DNA of said cells contain at least one FLP recombination target site therein.

15 In accordance with yet another aspect of the present invention, there are provided transgenic, non-human mammals, wherein said mammals contain at least one FLP recombination target site in the genomic DNA thereof.

20 In accordance with yet another aspect of the present invention, there is provided a method for the site-specific integration of transfected DNA into the genome of the above-described cells and/or transgenic, non-human mammals, said method comprising:

(i) contacting said genome with:

25 (a) FLP recombinase, and
(b) a first DNA comprising a nucleotide sequence containing at least one FLP recombination target site therein; and
30 thereafter

(ii) maintaining the product of Step (i) under conditions suitable for site-specific integration of said DNA sequence to occur at the FLP recombination target site in said genome.

35

-10-

In accordance with a further aspect of the present invention, there is provided a method for the analysis of the development of a mammal, said method comprising:

- 5 (a) providing a transgenic mammal comprising:
- (i) an expression construct encoding FLP under the control of a conditional promoter, and
 - 10 (ii) a reporter construct under the control of the same or a different promoter, wherein said reporter construct encodes a functional or non-functional reporter gene product, and wherein said construct contains
 - 15 at least one FLP recombination target site therein,
 - wherein the functional expression of the functional reporter gene is disrupted when said FLP recombination event occurs, or
 - 20 wherein the functional expression of the non-functional reporter gene commences when said FLP recombination event occurs; and
- 25 (b) following the development of said mammal to determine when expression of functional reporter gene product either commences or is disrupted.

In accordance with a still further aspect of the present invention, there is provided a co-transfection assay FLP-mediated recombination, said assay

30 comprising:

- (a) co-transfecting a host mammalian cell with:
- (i) a FLP expression plasmid, and
 - (ii) a reporter plasmid comprising a
 - 35 reporter gene inactivated by the presence

-11-

of at least one recombination target site;
and

- (b) monitoring said host cell under a variety of conditions for the gain of expression of functional reporter gene product.

FLP recombinase is a protein which catalyzes a site-specific recombination reaction that is involved in amplifying the copy number of the 2 μ plasmid of *S.cerevisiae* during DNA replication. FLP protein has been cloned and expressed in *E.coli* [see, for example, Cox, in proceedings of the National Academy of Sciences U.S.A., Vol. 80: 4223-4227 (1983)], and has been purified to near homogeneity [see, for example, Meyer-Lean, et al., in *Nucleic Acids Research*, Vol. 15: 6469-6488 (1987)]. FLP recombinases contemplated for use in the practice of the present invention are derived from species of the genus *Saccharomyces*. Preferred recombinases employed in the practice of the present invention are derived from strains of *Saccharomyces cerevisiae*. Especially preferred recombinases employed in the practice of the present invention are proteins having substantially the same amino acid sequence as set forth in Sequence I.D. No. 2, as encoded, for example, by Sequence I.D. No. 1, or the sequence set forth by Hartley and Donelson, *Nature* 286: 860 (1980).

The FLP recombination target site (sometimes referred to herein as "FRT") has also been identified as minimally comprising two 13 base-pair repeats, separated by an 8 base-pair spacer, as follows:

-Spacer-
5'-GAAGTTCCTATTC[TCTAGAAA]GTATAGGAACTTC-3'
XbaI
site

The nucleotides in the above "spacer" region can be replaced with any other combination of nucleotides, so

-12-

long as the two 13 base-pair repeats are separated by 8 nucleotides. The actual nucleotide sequence of the spacer is not critical, although those of skill in the art recognize that, for some applications, it is
5 desirable for the spacer to be asymmetric, while for other applications, a symmetrical spacer can be employed. Generally, the spacers of the FLP recombination target sites undergoing recombination with one another will be the same.

10 As schematically illustrated in Figure 1A, contact of genomic DNA containing a FLP recombination target site (shown as the linear Psv-BETA-GAL construct) with a vector containing a FLP recombination target site, in the presence of the protein, FLP recombinase, results
15 in recombination that forms a new genomic sequence wherein the vector sequences have been precisely incorporated into the genome of the host. The reverse of this process is shown schematically in Figure 1B, wherein a genomic sequence or construct containing two tandemly
20 oriented FLP recombination target sites, upon contacting with FLP, is recombined and the FLP recombination target site-bounded fragment is excised as a circular molecule.

Genes of interest contemplated for use in the practice of the present invention can be selected
25 from genes which provide a readily analyzable functional feature to the host cell and/or organism, e.g., visible markers (such as β -galactosidase, thymidine kinase, tyrosinase, and the like), selectable markers, (such as markers useful for positive and negative selection, e.g.,
30 genes for antibiotic resistance), as well as other functions which alter the phenotype of the recipient cells, and the like.

The first DNA employed in the practice of the present invention can comprise any nucleotide
35 sequence containing at least one FLP recombination target site, which will precisely define the locus at which FLP-

-13-

mediated recombination will occur. The nucleotide sequence can comprise all or part of a gene of interest, as well as other sequences not necessarily associated with any known gene. Optionally, for ease of later recovery of the gene of interest (in "activated" or modified form), the first DNA can optionally contain a second FLP recombination target site.

The second DNA employed in the practice of the present invention is selected from at least a second portion of the first gene of interest or at least a portion of a second gene of interest (including an intact form of a second gene of interest). When the second DNA is at least a second portion of the first gene of interest, the site-specific recombination of the present invention may act to provide a functional combination of the different portions of the first gene of interest. Alternatively, when the second DNA is at least a portion of a second gene of interest, the site-specific recombination of the present invention may act to provide a functional hybrid gene, which produces a product which is not identical with either the product of the first gene or the second gene. As yet another alternative, when the second DNA is a portion of a second gene, the site-specific recombination of the present invention may act to disrupt the function of the first gene of interest. Based on the nature of the first DNA and the second DNA, as well as the mode of interaction between the two, the site-specific interaction of the present invention may create or disrupt a feature which is colorimetrically detectable, immunologically detectable, genetically detectable, and the like.

In accordance with the present invention, assembly of a functional expression unit is achieved in any of a variety of ways, e.g., by association of the gene of interest with a functional promoter, by assembly of common gene fragments to produce a complete functional

-14-

gene (which, in combination with its promoter, comprises a functional expression unit), or assembly of diverse gene fragments from diverse sources to produce a functional, hybrid gene (which, in combination with a promoter, comprises a functional expression unit), and the like. Upon assembly of a functional expression unit as described herein, expression of the functional gene to produce a protein product can be activated in the usual manner. In the absence of FLP-mediated recombination, activation of expression would fail to produce a functional protein product.

In accordance with the present invention, dis-assembly of a functional expression unit is achieved in any of a variety of ways, e.g., by dis-associating the gene of interest from a functional promoter, by dis-assembly (e.g., disruption) of the functional gene (e.g., by introduction of DNA which renders the entire sequence non-functional), by removal of a substantial portion of the coding region of said gene, and the like. Upon dis-assembly of a functional expression unit as described herein, expression of the functional gene product under the conditions required prior to gene dis-assembly is no longer possible. The ability of the expression unit to be activated for expression has therefore been disrupted. The gene in this situation can be said to be inactivated, since activation of expression is not possible.

Individually inactive gene segments contemplated for use in the practice of the present invention are fragments which, alone, do not encode functional products. Such fragments can be derived from a first gene of interest alone, or from both a first and second gene of interest DNA fragments.

When gene inactivation is desired, the gene of interest can be disrupted with a DNA fragment which throws the gene of interest out of reading frame (e.g., an insert wherein the number of nucleotides is not

-15-

a multiple of 3). Alternatively, the gene of interest can be disrupted with a fragment which encodes a segment which is substantially dissimilar with the gene of interest so as to render the resulting product non-functional. As yet another alternative, the gene of interest can be disrupted so as to dis-associate the gene of interest from the transcriptional control of the promoter with which it is normally associated.

The introduction of DNA, e.g., DNA encoding FLP recombination target sites, into the genome of target cells can be accomplished employing standard techniques, e.g., transfection, microinjection, electroporation, infection with retroviral vectors, and the like.

Introduction of protein, e.g., FLP recombinase protein, to host cells and/or organisms can be accomplished by standard techniques, such as for example, injection or microinjection, transfection with nucleotide sequences encoding FLP, and the like.

When employed for gene therapy of an intact organism, introduction of transgenic cells into the subject is accomplished by standard techniques, such as for example, grafting, implantation, and the like.

Mammalian cells contemplated for use in the practice of the present invention include all members of the order Mammalia, such as, for example, human cells, mouse cells, rat cells, monkey cells, hamster cells, and the like.

Host organisms contemplated for use in the practice of the present invention include each of the organism types mentioned above, with the proviso, however, that no claim is made to genetically modified human hosts (although the present invention contemplates methods for the treatment of humans).

Once FLP recombinase (or DNA encoding same) and DNA containing at least one FLP recombination

-16-

target site have been introduced into suitable host cells/organisms, the cells/host organisms are maintained under conditions suitable for the site-specific recombination of DNA. Such conditions generally involve conditions required for the viability of the host cell or organism. For in vitro manipulations, conditions employed typically involve low concentrations of a variety of buffers having a pH of between about 5-9 and ionic strengths in the range of about 50-350 mM. See, for example, Senecoff, et al., in Journal of Molecular Biology, Vol. 201: 405-421 (1988).

In accordance with a particular aspect of the present invention, a co-transfection assay has been developed which can be used to characterize FLP-mediated recombination of extrachromosomal DNA in a variety of cell lines. Cells are co-transfected with an expression construct and a "reporter" plasmid that is a substrate for the recombinase. The expression construct encodes a FLP recombinase protein. The reporter plasmid encodes either a functional reporter gene containing at least one recombination target site therein, or a non-functional reporter gene containing at least one recombination target site therein. Upon expression of FLP by the expression construct, the functional reporter gene will be rendered non-functional, or the non-functional reporter gene will be rendered functional. Thus, the activity of the expression construct can be assayed either by recovering the reporter plasmid and looking for evidence of recombination at the DNA level, or by preparing cytoplasmic extracts and looking for evidence of recombination at the protein level (i.e., by measuring the expression of reporter gene activity generated by the recombined reporter). Such assays are described in greater detail in Example 1 below.

-17-

The invention will now be described in greater detail by reference to the following non-limiting examples.

5

EXAMPLES

Example 1

Co-transfection Assays

10 The co-transfection assay used to characterize FLP-mediated recombination of extrachromosomal DNA involved transfection of cells with an expression construct and a "reporter" plasmid that was a substrate for the recombinase. The activity of the
15 expression construct could be assayed either by recovering the reporter plasmid and looking for molecular evidence of recombination at the DNA level, or by preparing cytoplasmic extracts and looking for evidence of recombination at the protein level (i.e., by measuring
20 β -galactosidase activity generated by recombined reporter).

 The pNEO β GAL reporter plasmid used for these assays was derived from pFRT β GAL (Fig. 2A). In the Figure, half-arrows indicate positions of FLP
25 recombination target (FRT) sites; E and S designate EcoRI and ScaI restriction sites, respectively; Psv designates early promoter from SV40; BETA-GAL designates the β -galactosidase structural sequence; NEO designates neomycin expression cassette; Pcmv designates the
30 cytomegalovirus immediate early promoter; IN designates an intron; FLP designates a FLP coding sequence; AN designates an SV40 adenylation cassette; thin lines represent vector sequences; and the sizes of restriction fragments are indicated in kb.

35 pFRT β GAL contains a version of the bacterial β -galactosidase sequence modified by insertion

-18-

of a FLP recombination target site, or FRT, within the protein coding sequence immediately 3' to the translational start. The oligonucleotide used for the construction of pFT β RGAL was:

5

5'-GATCCCGGGCTACCATGGA•GAAGTTCCTATTC•CGAAGTTCCTATTC
(TCTAGA)AAGTATAGGAACTTCA-3'.

This oligonucleotide contains an in-frame start codon, minimal FRT site, and an additional copy of the 13-bp FRT repeat [•XXX•]; the XbaI site within the FRT spacer is enclosed in parentheses. The linker was inserted between the BamHI and HindIII sites of pSKS105 (M.J. Casadaban, A. Martin-Arias, S.K. Shapira, and J. Chou, *Meth. Enzymol.* 100, 293 (1983)) and the LacZ portion of modified gene was cloned into a pSV2 vector. The neomycin cassette used for construction of pNEO β GAL was an XhoI to BamHI fragment from pMC1neo-polyA (K. Thomas and M. Capecchi, *Cell* 51:503 (1987)) cloned between copies of the J3 FRT site in pUC19.

The FRT consists of two inverted 13-base-pair (bp) repeats and an 8-bp spacer that together comprise the minimal FRT site, plus an additional 13-bp repeat which may augment reactivity of the minimal substrate. The β -galactosidase translational reading frame was preserved upon insertion of the FRT site, and the resulting plasmid, pFRT β GAL, generated robust activity in mammalian cells (Table 1).

pNEO β GAL was constructed by cutting pFRT β GAL in the middle of the FRT site with XbaI and then inserting an XbaI fragment consisting of two half-FRT sites flanking a neomycin transcription unit. This created intact FRTs on either side of the neomycin cassette and rendered the β -galactosidase transcription unit inactive (Table 1). Precise FLP-mediated recombination of the FRTs caused the excision of the

-19-

neomycin cassette, recreated the parental pFRT β GAL plasmid, and restored β -galactosidase expression.

Co-transfection of cells with a fixed amount of pNEO β GAL reporter plasmid and increasing amounts of the pOG44 FLP expression vector generated increasing amounts of recombined reporter plasmid and consequently, increased levels of β -galactosidase activity. Molecular evidence for FLP-mediated recombination was obtained by recovering plasmids 36 hours after transfection, followed by endonuclease treatment (with EcoRI and ScaI) and Southern blotting (see Fig. 2B; employing as a probe the fragment of pFRT β GAL indicated at the top of Fig. 2A). Lysates of cells from cotransfections that included the pOG44 FLP expression vector showed a signal at 5.6 kb, the position at which recombined reporter (equivalent to pFRT β GAL) would run, and a 3.2 kb signal that was generated by unrecombined pNEO β GAL reporter (Fig. 2A). The 5.6 kb band intensity was proportional to the amount of FLP expression plasmid included in the transfection. The 5.6 kb band was not seen in cotransfections in which a non-FLP plasmid was substituted for the FLP expression vector (Fig. 2B) or in transfections that contained only pOG44 (and no reporter plasmid). pOG44 generated additional signals at 2.2 kb and 2.8 kb because the plasmid used in its construction contained EcoRI and EcoRI-ScaI fragments of such length.

pOG44 consists of the cytomegalovirus immediate early promoter from pCDM8 [see Aruffo and Seed in Proc. Natl Acad. Sci. , USA 84:8573 (1987)], a 5' leader sequence and synthetic intron from pMLSIScat [see Huang and Gorman in Nucl. Acids Res. 18: 937 (1990)], the FLP coding sequence (bp 5568-6318 and 1-626 of the 2 μ m circle, [see Hartley and Donelson, Nature 286: 860 (1980)] and the SV40 late region polyadenylation signal from pMLSIScat. The following silent nucleotide

-20-

substitutions were introduced into the structural FLP sequence using the polymerase chain reaction: C for T at position 5791, G for A at 5794, G for C at 5800, C for T at 55, G for A at 58, and C for T at 103. These changes eliminated three canonical AATAAA polyadenylation signals and introduced a PstI restriction site without altering the amino acid sequence encoded by the nucleotide sequence. pOG28 consists of a murine cDNA for dihydrofolate reductase cloned into pCDM8 (Aruffo and Seed, supra).

In the same samples, β -galactosidase activity was also proportional to the amount of FLP expression plasmid included (Fig. 2C). Only background activities were observed in cotransfections that included a non-FLP control plasmid (Table 1) or when pOG44 alone was transfected. The experiment thus provides both molecular and biochemical evidence for precise FLP-mediated recombination in mammalian cells.

Table 1 presents β -galactosidase activities in cotransfection assays of 293, CV-1, and F-9 cells. Positive control transfections (pFRT β GAL) included 1 μ g of pFRT β GAL and 18 μ g of the pOG28 non-FLP control plasmid; negative control transfections (pNEO β GAL) included 1 μ g of pNEO β GAL and 18 μ g of the pOG28; and experimental transfections (pNEO β GAL + FLP) contained 1 μ g of pNEO β GAL and 18 μ g of the pOG44 FLP expression plasmid (Fig. 1A). The column headed by "%" shows the pNEO β GAL + FLP values as a percentage of the pFRT β GAL positive control. Each value represents the mean for six plates from two experiments. Standard errors are in parentheses. Neither pOG28 nor pOG44 generated β -galactosidase activity when transfected alone. All transfections contained 1 μ g of pRSVL [de Wet et al., Mol. Cell. Biol. 7: 725 (1987)] to correct β -galactosidase activities for relative transfection efficiencies.

-21-

Subconfluent cultures of cells in 10 cm dishes and grown in Dulbecco's modified Eagle's medium (DMEM) and 5% calf serum were transfected by overnight exposure to calcium phosphate precipitates [Graham et al., Virology 36:59 (1979)] and then split 1:4. After 24 hours incubation, one plate of each transfection was harvested by Hirt extraction [J. Mol Biol. 26:365 (1967)] and a second plate was used to prepare cytoplasmic extracts [de Wet et al., supra]. Approximately 5% of the DNA recovered from single plates was used for Southern analyses. β -galactosidase assays were performed as described by Hall et al., in J. Mol. Appl. Genet. 2:101 (1983)]. Luciferase activities generated by the inclusion of 1 μ g of pRSVL (de Wet et al., supra) in all transfections were used to correct β -galactosidase activities for relative transfection efficiencies. The experiment was repeated twice with similar results.

TABLE 1: β -GALACTOSIDASE ACTIVITIES (UNITS/ MG PROTEIN) IN COTRANSFECTED CELLS

CELL LINE	TRANSFECTIONS			%
	pFRT β GAL	pNEO β GAL	pNEO β GAL + FLP	
293	30.4 (1.9)	0.17 (0.02)	14.2 (2.2)	47
CV-1	275 (25)	0.33 (0.06)	22.6 (1.2)	8
F9	24.8 (4.3)	0.04 (0.01)	1.88 (0.02)	8

FLP activity has also been demonstrated in monkey kidney (CV-1) and mouse embryonal carcinoma (F9) cells. In Table 1, the β -galactosidase activity in the "pFRT β GAL" transfections represents an estimate of the expression expected if all the pNEO β GAL in a co-

-22-

transfection were immediately recombined. The highest β -galactosidase expression in co-transfections employing pNEO β GAL plus pOG44, relative to pFRT β GAL transfected cells, was 47%, seen in 293 cells. This is a remarkable level considering that β -galactosidase expression required both FLP expression, followed by recombination of pNEO β GAL, to produce a construct capable of expressing β -galactosidase. Co-transfections of CV-1 and F9 cells generated 8% of the activity seen in the pFRT β GAL transfections. Even at this lower relative activity, cotransfected cells were readily observed in histochemical reactions for β -galactosidase activity.

Example 2

FLP-Mediated Removal of Intervening Sequences

If the invention method is to be widely applicable, for example for gene activation in transgenic mammals, the ability of FLP to faithfully promote precise recombination at FLP recombination target sites contained in the mammalian genome is required. Such ability is demonstrated in this example.

Cell lines that contain single integrated copies of pNEO β GAL (designated CVNEO β GAL/E) were produced by transfecting CV-1 cells with linearized plasmid by electroporation, then isolated by selecting G418-resistant (G418^R) transfectants that stably expressed the neomycin cassette, and finally identifying single copy lines by Southern blot analyses (Fig. 3). As previously shown for other integrated constructs with similarly short direct repeats, the chromosomal FRTs did not spontaneously recombine (in the absence of FLP) to produce a β -galactosidase-positive (β GAL⁺) phenotype at detectable frequencies (Table 2).

-23-

Transient expression of FLP in the CVNEO β GAL/E lines (by transiently transfecting with the pOG44 FLP expression vector) promoted a rapid conversion to a β GAL⁺ phenotype. When five different lines were

5 transiently transfected with the pOG44 FLP expression vector, β -galactosidase activities at 36 hours were 40 to 100-fold higher than those seen in replicate plates transfected with a non-FLP plasmid (Table 2). At 48

10 hours after transfection histochemical processing showed many positive cells (Table 2).

Table 2 presents the β -galactosidase phenotypes of CVNEO β GAL/E lines, which contain a single copy of the β -galactosidase inactive reporter, pNEO β GAL, after transfection with FLP expression (pOG44), non-FLP

15 negative control (pOG28) or β -galactosidase positive control (pFRT β GAL) plasmids. The pFRT β GAL transfections included 1 μ g of pFRT β GAL and 19 μ g of pOG44; other mixes contained 20 μ g of the indicated plasmid. β -galactosidase

20 activities are mean values for triplicate transfections performed as described for Fig. 2 and assayed 36 hours after removal of precipitates; standard errors for the pOG44 transfections were less than 10% of the mean. The percent positive was determined by scoring more than 10³ cells after transfection and histochemical processing as

25 described by de Wet et al., supra.

-24-

TABLE 2

β -GALACTOSIDASE PHENOTYPES OF
TRANSFECTED CVNEO β GAL CELL LINES

5	CELL LINE	ACTIVITIES (units/mg protein)		PERCENT POSITIVE		
		pOG28	pOG44	pOG28	pFRT β GAL	pOG44
10	E6	0.24	11.2	0†	8.7	6.1
	E25	0.21	16.7	0†	17.1	12.4
15	E26	0.18	7.2	0†	19.5	15.4
	E14	0.28	13.1	ND	ND	ND
	E22	0.09	9.6	ND	ND	ND

†No positive cells were found among $>10^6$ cells examined.

20 ND: Not done.

To provide some estimate of the efficiency of recombination, an additional set of replicate plates were transfected with the pFRT β GAL β -galactosidase expression vector. Comparing the fractions of cells that were β GAL⁺ in the pFRT β GAL and in the pOG44 transfections (assuming similar transfection efficiencies) suggests that most (70-80%) of the cells transfected with pOG44 were converted to a β GAL⁺ phenotype (Table 2). The comparison undoubtedly underestimates the efficiency of FLP-mediated excision. Whereas many copies of a functional β -galactosidase gene were available for immediate transcription in the positive controls, recombination may have occurred shortly before harvest in some pOG44-transfected cells. In these cases the single recombined reporter gene may not have generated enough β -galactosidase by the time of harvest to render the cells positive in this assay.

-25-

The β GAL⁺ phenotype was passed on to all descendents of many FLP-converted cells. Positive colonies were formed during prolonged expansion of individual colonies. Entirely negative colonies and mixed colonies were also observed. Mixed colonies would be expected if recombination occurred after mitosis in only one descendent of a transfected cell, or if recombined and unrecombined cells mixed at replating or during subsequent growth. Indeed, the physical segregation of phenotypes evident in most mixed colonies suggested that they were composed of stably positive and negative lineages.

The correlation between β -galactosidase expression and recombination at FRT sites was examined by comparing the structure of the integrated pNEO β GAL sequences in two β GAL⁺ subclones to the parental line. CVNEO β GAL/E25 (106) cells were transfected with the pOG44 FLP expression vector and subcloned 12 hours after removal of the precipitate. After histochemical screening, two β GAL⁺ subclones (E25B1 and E25B2) were expanded for further analysis. In Southern blots of genomic DNA from both subclones, the pattern of hybridization matched that expected for FLP-mediated recombination of the FRT sites in the parental line (Fig. 3). While recombination products have not been recovered and sequenced, these Southern analyses and the fact that activation of β -galactosidase expression required creation of a functional translational reading frame indicate that FLP-mediated recombination was precise.

30

-26-

Example 3

FLP Mediated Recombination of FRT
on an Extrachromosomal Molecule With a
Chromosomally Integrated FRT

5

Reversal of the process described in the previous Example, i.e., the FLP-mediated recombination of an FRT site on a plasmid with a chromosomally integrated FRT site, can be used to target the integration of transfected plasmids to specific genomic sites. To determine the frequency at which this occurs, G418-sensitive, β GAL⁺ E25B2 cells were co-transfected with the pOG44 FLP expression vector and a plasmid, pOG45, that contained a neomycin resistance gene expression cassette and a single FRT. pOG45 consisted of the neomycin resistance cassette and 3' FRT from pNEO β GAL cloned into pUC19. 8×10^5 CVNEO β GAL cells were transfected by electroporation in 800 μ l of saline containing 40 μ g of pOG44 and 0.1 μ g of either pOG45 or, for a negative control, pOG45A (which was derived from pOG45 by deleting a 200 bp fragment containing the FRT).

G418^R subclones (designated B2N) from three transfections that had stably integrated pOG45 were histochemically stained for β -galactosidase activity and more than half (104 of 158, or 66%) were either entirely β -galactosidase-negative (β GAL⁻) or predominantly β GAL⁻ with a few clusters of β GAL⁺ cells. The remaining colonies were β GAL⁺. With continued passage as dispersed monolayers, the fraction of β GAL⁺ cells in the mosaic lines rapidly diminished. This suggested they were G418-sensitive cells that initially survived because of their proximity to resistant cells; this was confirmed by reconstitution experiments. All of the 55 colonies formed after parallel co-transfections of pOG44 and a derivative of pOG45 (pOG45A) that lacked an FRT were β GAL⁺.

-27-

The correlation between loss of β -galactosidase activity and recombination between plasmid and chromosomal FRTs was examined in Southern analyses. Because the FRT and neomycin cassette of pOG45 were derived from the neomycin cassette and 3' FRT of pNEO β GAL (Fig. 2A), recombination of the plasmid FRT with the E25B2 chromosomal FRT regenerates the 3.2 kb EcoRI fragment of the original CVNEO β GAL/E25 parent. Additionally, the 8.5 kb junctional fragment of CVNEO β GAL/E25 shifts to 12.0 kb because pOG45 is 3.5 kb larger than the neomycin cassette of pNEO β GAL. The 3.2 kb EcoRI fragment and the 8.5 kb junctional fragment were observed in each of the 10 cell lines analyzed after initial histochemical classification as β GAL⁻ or mosaic, as shown for two such lines in Fig. 3B. In contrast, each of the four β GAL⁺ colonies examined by Southern analyses showed that pOG45 had integrated at a random site.

These data show that FLP-mediated recombination will target the integration of transfected DNA to a specific chromosomal site at frequencies that exceed those of random integration, and that the event can be marked by the alteration in gene activity at the target site. The efficiency of targeted integration can be increased by standard optimization techniques, such as for example, by using ratios of the integrating plasmid and FLP expression vectors different from the single ratio mixture used here, or by using FRT mutations in the plasmid and chromosomal sites to decrease the frequency with which successfully integrated plasmids are subsequently excised.

While the invention has been described in detail with reference to certain preferred embodiments thereof, it will be understood that modifications and variations are within the spirit and scope of that which is described and claimed.

-28-

SUMMARY OF SEQUENCES

Sequence I.D. No. 1 is the approximately
1450 base-pair sequence encoding a FLP recombinase
5 contemplated for use in the practice of the present
invention, as well as the amino acid sequence deduced
therefrom.

Sequence I.D. No. 2 is the amino acid
sequence deduced from the nucleotide sequence of Sequence
10 ID No. 1.

-29-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: WAHL, DR, GEOFFREY M
O'GORMAN DR, STEPHEN V
- (ii) TITLE OF INVENTION: FLP-MEDIATED GENE MODIFICATION IN
MAMMALIAN CELLS, AND COMPOSITIONS AND CELLS USEFUL
THEREFOR
- (iii) NUMBER OF SEQUENCES: 2
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: FITCH, EVEN, TABIN & FLANNERY
 - (B) STREET: 135 South LaSalle Street, Suite 510
 - (C) CITY: Chicago
 - (D) STATE: Illinois
 - (E) COUNTRY: USA
 - (F) ZIP: 60603
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: US
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: REITER MR, STEPHEN E
 - (B) REGISTRATION NUMBER: 31192
 - (C) REFERENCE/DOCKET NUMBER: 50730
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (619) 552-1311
 - (B) TELEFAX: (619) 552-0095
 - (C) TELEX: 20 6566 PATLAW CGO

-30-

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1380 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vii) IMMEDIATE SOURCE:

(B) CLONE: NATIVE FLP

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..1269

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG	CCA	CAA	TTT	GAT	ATA	TTA	TGT	AAA	ACA	CCA	CCT	AAG	GTG	CTT	GTT	48
Met	Pro	Gln	Phe	Asp	Ile	Leu	Cys	Lys	Thr	Pro	Pro	Lys	Val	Leu	Val	
1				5					10					15		
CGT	CAG	TTT	GTG	GAA	AGG	TTT	GAA	AGA	CCT	TCA	GGT	GAG	AAA	ATA	GCA	96
Arg	Gln	Phe	Val	Glu	Arg	Phe	Glu	Arg	Pro	Ser	Gly	Glu	Lys	Ile	Ala	
			20					25					30			
TTA	TGT	GCT	GCT	GAA	CTA	ACC	TAT	TTA	TGT	TGG	ATG	ATT	ACA	CAT	AAC	144
Leu	Cys	Ala	Ala	Glu	Leu	Thr	Tyr	Leu	Cys	Trp	Met	Ile	Thr	His	Asn	
			35				40					45				
GGA	ACA	GCA	ATC	AAG	AGA	GCC	ACA	TTC	ATG	AGC	TAT	AAT	ACT	ATC	ATA	192
Gly	Thr	Ala	Ile	Lys	Arg	Ala	Thr	Phe	Met	Ser	Tyr	Asn	Thr	Ile	Ile	
			50			55					60					
AGC	AAT	TCG	CTG	AGT	TTC	GAT	ATT	GTC	AAT	AAA	TCA	CTC	CAG	TTT	AAA	240
Ser	Asn	Ser	Leu	Ser	Phe	Asp	Ile	Val	Asn	Lys	Ser	Leu	Gln	Phe	Lys	
65					70					75					80	
TAC	AAG	ACG	CAA	AAA	GCA	ACA	ATT	CTG	GAA	GCC	TCA	TTA	AAG	AAA	TTG	288
Tyr	Lys	Thr	Gln	Lys	Ala	Thr	Ile	Leu	Glu	Ala	Ser	Leu	Lys	Lys	Leu	
				85					90					95		
ATT	CCT	GCT	TGG	GAA	TTT	ACA	ATT	ATT	CCT	TAC	TAT	GGA	CAA	AAA	CAT	336
Ile	Pro	Ala	Trp	Glu	Phe	Thr	Ile	Ile	Pro	Tyr	Tyr	Gly	Gln	Lys	His	
			100					105					110			
CAA	TCT	GAT	ATC	ACT	GAT	ATT	GTA	AGT	AGT	TTG	CAA	TTA	CAG	TTC	GAA	384
Gln	Ser	Asp	Ile	Thr	Asp	Ile	Val	Ser	Ser	Leu	Gln	Leu	Gln	Phe	Glu	
			115				120					125				
TCA	TCG	GAA	GAA	GCA	GAT	AAG	GGA	AAT	AGC	CAC	AGT	AAA	AAA	ATG	CTT	432
Ser	Ser	Glu	Glu	Ala	Asp	Lys	Gly	Asn	Ser	His	Ser	Lys	Lys	Met	Leu	
			130			135					140					
AAA	GCA	CTT	CTA	AGT	GAG	GGT	GAA	AGC	ATC	TGG	GAG	ATC	ACT	GAG	AAA	480
Lys	Ala	Leu	Leu	Ser	Glu	Gly	Glu	Ser	Ile	Trp	Glu	Ile	Thr	Glu	Lys	
145					150					155					160	

-31-

ATA	CTA	AAT	TCG	TTT	GAG	TAT	ACT	TCG	AGA	TTT	ACA	AAA	ACA	AAA	ACT	528
Ile	Leu	Asn	Ser	Phe	Glu	Tyr	Thr	Ser	Arg	Phe	Thr	Lys	Thr	Lys	Thr	
				165					170					175		
TTA	TAC	CAA	TTC	CTC	TTC	CTA	GCT	ACT	TTC	ATC	AAT	TGT	GGA	AGA	TTC	576
Leu	Tyr	Gln	Phe	Leu	Phe	Leu	Ala	Thr	Phe	Ile	Asn	Cys	Gly	Arg	Phe	
			180					185					190			
AGC	GAT	ATT	AAG	AAC	GTT	GAT	CCG	AAA	TCA	TTT	AAA	TTA	GTC	CAA	AAT	624
Ser	Asp	Ile	Lys	Asn	Val	Asp	Pro	Lys	Ser	Phe	Lys	Leu	Val	Gln	Asn	
		195					200					205				
AAG	TAT	CTG	GGA	GTA	ATA	ATC	CAG	TGT	TTA	GTG	ACA	GAG	ACA	AAG	ACA	672
Lys	Tyr	Leu	Gly	Val	Ile	Ile	Gln	Cys	Leu	Val	Thr	Glu	Thr	Lys	Thr	
	210					215					220					
AGC	GTT	AGT	AGG	CAC	ATA	TAC	TTC	TTT	AGC	GCA	AGG	GGT	AGG	ATC	GAT	720
Ser	Val	Ser	Arg	His	Ile	Tyr	Phe	Phe	Ser	Ala	Arg	Gly	Arg	Ile	Asp	
					230					235					240	
CCA	CTT	GTA	TAT	TTG	GAT	GAA	TTT	TTG	AGG	AAT	TCT	GAA	CCA	GTC	CTA	768
Pro	Leu	Val	Tyr	Leu	Asp	Glu	Phe	Leu	Arg	Asn	Ser	Glu	Pro	Val	Leu	
				245					250					255		
AAA	CGA	GTA	AAT	AGG	ACC	GGC	AAT	TCT	TCA	AGC	AAT	AAA	CAG	GAA	TAC	816
Lys	Arg	Val	Asn	Arg	Thr	Gly	Asn	Ser	Ser	Ser	Asn	Lys	Gln	Glu	Tyr	
			260					265					270			
CAA	TTA	TTA	AAA	GAT	AAC	TTA	GTC	AGA	TCG	TAC	AAT	AAA	GCT	TTG	AAG	864
Gln	Leu	Leu	Lys	Asp	Asn	Leu	Val	Arg	Ser	Tyr	Asn	Lys	Ala	Leu	Lys	
		275					280					285				
AAA	AAT	GCG	CCT	TAT	TCA	ATC	TTT	GCT	ATA	AAA	AAT	GCG	CCA	AAA	TCT	912
Lys	Asn	Ala	Pro	Tyr	Ser	Ile	Phe	Ala	Ile	Lys	Asn	Gly	Pro	Lys	Ser	
		290				295					300					
CAC	ATT	GGA	AGA	CAT	TTG	ATG	ACC	TCA	TTT	CTT	TCA	ATG	AAG	GGC	CTA	960
His	Ile	Gly	Arg	His	Leu	Met	Thr	Ser	Phe	Leu	Ser	Met	Lys	Gly	Leu	
	305				310					315					320	
ACG	GAG	TTG	ACT	AAT	GTT	GTG	GGA	AAT	TGG	AGC	GAT	AAG	CGT	GCT	TCT	1008
Thr	Glu	Leu	Thr	Asn	Val	Val	Gly	Asn	Trp	Ser	Asp	Lys	Arg	Ala	Ser	
				325					330					335		
GCC	GTG	GCC	AGG	ACA	ACG	TAT	ACT	GAT	CAG	ATA	ACA	GCA	ATA	CCT	GAT	1056
Ala	Val	Ala	Arg	Thr	Thr	Tyr	Thr	His	Gln	Ile	Thr	Ala	Ile	Pro	Asp	
			340					345					350			
CAC	TAC	TTC	GCA	CTA	GTT	TCT	CGG	TAC	TAT	GCA	TAT	GAT	CCA	ATA	TCA	1104
His	Tyr	Phe	Ala	Leu	Val	Ser	Arg	Tyr	Tyr	Ala	Tyr	Asp	Pro	Ile	Ser	
		355					360					365				
AAG	GAA	ATG	ATA	GCA	TTG	AAG	GAT	GAG	ACT	AAT	CCA	ATT	GAG	GAG	TGG	1152
Lys	Glu	Met	Ile	Ala	Leu	Lys	Asp	Glu	Thr	Asn	Pro	Ile	Glu	Glu	Trp	
		370				375					380					
CAG	CAT	ATA	GAA	CAG	CTA	AAG	GGT	AGT	GCT	GAA	GGA	AGC	ATA	CGA	TAC	1200
Gln	His	Ile	Glu	Gln	Leu	Lys	Gly	Ser	Ala	Glu	Gly	Ser	Ile	Arg	Tyr	
					390					395					400	

-32-

CCC GCA TGG AAT GGG ATA ATA TCA CAG GAG GTA CTA GAC TAC CTT TCA 1248
 Pro Ala Trp Asn Gly Ile Ile Ser Gln Glu Val Leu Asp Tyr Leu Ser
 405 410 415
 TCG TAC ATA AAT AGA CGC ATA TAAGTACGCA TTTAAGCATA AACACGCACT 1299
 Ser Tyr Ile Asn Arg Arg Ile
 420
 ATGCCGTTCT TCTCATGTAT ATATATATAC AGGCAACACG CAGATATAGG TGCGACGTGA 1359
 ACAGTGAGCT GTATGTGCGC A 1380

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 423 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Pro Gln Phe Asp Ile Leu Cys Lys Thr Pro Pro Lys Val Leu Val
 1 5 10 15
 Arg Gln Phe Val Glu Arg Phe Glu Arg Pro Ser Gly Glu Lys Ile Ala
 20 25 30
 Leu Cys Ala Ala Glu Leu Thr Tyr Leu Cys Trp Met Ile Thr His Asn
 35 40 45
 Gly Thr Ala Ile Lys Arg Ala Thr Phe Met Ser Tyr Asn Thr Ile Ile
 50 55 60
 Ser Asn Ser Leu Ser Phe Asp Ile Val Asn Lys Ser Leu Gln Phe Lys
 65 70 75 80
 Tyr Lys Thr Gln Lys Ala Thr Ile Leu Glu Ala Ser Leu Lys Lys Leu
 85 90 95
 Ile Pro Ala Trp Glu Phe Thr Ile Ile Pro Tyr Tyr Gly Gln Lys His
 100 105 110
 Gln Ser Asp Ile Thr Asp Ile Val Ser Ser Leu Gln Leu Gln Phe Glu
 115 120 125
 Ser Ser Glu Glu Ala Asp Lys Gly Asn Ser His Ser Lys Lys Met Leu
 130 135 140
 Lys Ala Leu Leu Ser Glu Gly Glu Ser Ile Trp Glu Ile Thr Glu Lys
 145 150 155 160
 Ile Leu Asn Ser Phe Glu Tyr Thr Ser Arg Phe Thr Lys Thr Lys Thr
 165 170 175
 Leu Tyr Gln Phe Leu Phe Leu Ala Thr Phe Ile Asn Cys Gly Arg Phe
 180 185 190

-33-

Ser Asp Ile Lys Asn Val Asp Pro Lys Ser Phe Lys Leu Val Gln Asn
 195 200 205
 Lys Tyr Leu Gly Val Ile Ile Gln Cys Leu Val Thr Glu Thr Lys Thr
 210 215 220
 Ser Val Ser Arg His Ile Tyr Phe Phe Ser Ala Arg Gly Arg Ile Asp
 225 230 235 240
 Pro Leu Val Tyr Leu Asp Glu Phe Leu Arg Asn Ser Glu Pro Val Leu
 245 250 255
 Lys Arg Val Asn Arg Thr Gly Asn Ser Ser Ser Asn Lys Gln Glu Tyr
 260 265 270
 Gln Leu Leu Lys Asp Asn Leu Val Arg Ser Tyr Asn Lys Ala Leu Lys
 275 280 285
 Lys Asn Ala Pro Tyr Ser Ile Phe Ala Ile Lys Asn Gly Pro Lys Ser
 290 295 300
 His Ile Gly Arg His Leu Met Thr Ser Phe Leu Ser Met Lys Gly Leu
 305 310 315 320
 Thr Glu Leu Thr Asn Val Val Gly Asn Trp Ser Asp Lys Arg Ala Ser
 325 330 335
 Ala Val Ala Arg Thr Thr Tyr Thr His Gln Ile Thr Ala Ile Pro Asp
 340 345 350
 His Tyr Phe Ala Leu Val Ser Arg Tyr Tyr Ala Tyr Asp Pro Ile Ser
 355 360 365
 Lys Glu Met Ile Ala Leu Lys Asp Glu Thr Asn Pro Ile Glu Glu Trp
 370 375 380
 Gln His Ile Glu Gln Leu Lys Gly Ser Ala Glu Gly Ser Ile Arg Tyr
 385 390 395 400
 Pro Ala Trp Asn Gly Ile Ile Ser Gln Glu Val Leu Asp Tyr Leu Ser
 405 410 415
 Ser Tyr Ile Asn Arg Arg Ile
 420

-34-

THAT WHICH IS CLAIMED IS:

1. A mammalian recombination system comprising:
 - 5 (i) FLP recombinase, or a nucleotide sequence encoding same, and
 - (ii) a first DNA comprising a nucleotide sequence containing at least one FLP recombination target site therein.
- 10 2. A recombination system according to Claim 1 further comprising:
 - (iii) a second DNA, wherein said second DNA is selected from:
 - 15 (a) at least a second portion of said first gene of interest, or
 - (b) at least a portion of a second gene of interest;wherein said second DNA contains at least one FLP
20 recombination target site; and wherein said second DNA, when combined in reading frame with said first DNA, provides a functional gene.
- 25 3. A recombination system according to Claim 2 wherein said second DNA comprises an additional portion of said first gene of interest.
- 30 4. A recombination system according to Claim 2 wherein said second DNA comprises at least a portion of a second gene of interest.
5. A recombination system according to Claim 4 wherein said portion of said second gene of interest,

-35-

when combined in reading frame with said first DNA,
provides a hybrid, functional gene.

6. A recombination system according to Claim 4
5 wherein said portion of said second gene of interest,
when combined with said first DNA, disrupts the function
of said first gene of interest.

7. A recombination system according to Claim 1
10 wherein said first DNA further comprises a second FLP
recombination target site.

8. A recombination system according to Claim 1
wherein the FLP recombinase is derived from a species of
15 the genus *Saccharomyces*.

9. A recombination system according to Claim 1
wherein the FLP recombinase is derived from a strain of
Saccharomyces cerevisiae.
20

10. A recombination system according to Claim 9
wherein said FLP recombinase is encoded by the
approximately 1450 base pair sequence set forth as
Sequence ID No. 1.
25

11. A recombination system according to Claim 1
wherein said first DNA provides a readily analyzable
marker feature to the host system.

12. A recombination system according to Claim 2
30 wherein said second DNA provides a readily analyzable
marker feature to the host system.

-36-

13. A DNA construct comprising, as an autonomous fragment:

- (a) at least one FLP recombination target site,
- 5 (b) at least one restriction endonuclease recognition site,
- (c) at least one marker gene,
- (d) a bacterial origin of replication, and optionally
- 10 (e) a mammalian cellular or viral origin of DNA replication.

14. A DNA construct comprising, as an autonomous fragment, in the following order, reading from
15 5' to 3' along said fragment:

- (a) a first FLP recombination target site,
- (b) an insert portion comprising, in any suitable sequence:
 - (1) at least one restriction
 - 20 endonuclease recognition site,
 - (2) at least one marker gene,
 - (3) a bacterial origin of replication, and optionally
 - (4) a mammalian cellular or viral origin
 - 25 of DNA replication, and
- (c) a second FLP recombination target site in tandem with said first FLP recombination target site.

30 15. A method for the assembly of functional gene(s), which is (are) then suitable for activation of expression in mammalian cells, by recombination of individually inactive gene segments derived from one or

-37-

more gene(s) of interest, wherein each of said segments contains at least one recombination target site, said method comprising:

5 contacting said individually inactive gene segments with a FLP recombinase, under conditions suitable for recombination to occur, thereby providing a DNA sequence which encodes a functional gene of interest.

10 16. A method according to Claim 15 wherein the FLP recombinase is derived from a species of the genus *Saccharomyces*.

15 17. A method according to Claim 15 wherein the FLP recombinase is derived from a strain of *Saccharomyces cerevisiae*.

20 18. A method according to Claim 17 wherein said FLP recombinase is encoded by the approximately 1450 base pair sequence set forth as Sequence ID No. 1.

25 19. A method for the disruption of functional gene(s) of interest, rendering said gene(s) unable to be inactivated for expression in mammalian cells wherein said gene(s) of interest contain at least one FLP recombination target site, said method comprising contacting said gene(s) of interest with:

30 (i) a DNA segment which contains at least one FLP recombination target site, and
 (ii) FLP recombinase;

wherein said contacting is carried out under conditions suitable for recombination to occur between said gene and

-38-

said DNA segment, thereby disrupting the gene(s) of interest and rendering said gene(s) non-functional.

20. A method according to Claim 19 wherein said
5 DNA segment provides a readily analyzable marker feature to the host system.

21. A method according to Claim 19 wherein the
FLP recombinase is derived from a species of the genus
10 *Saccharomyces*.

22. A method according to Claim 19 wherein the
FLP recombinase is derived from a strain of *Saccharomyces cerevisiae*.
15

23. A method according to Claim 22 wherein said
FLP recombinase is encoded by the approximately 1450 base
pair sequence set forth as Sequence ID No. 1.

20 24. A method for the recovery of transfected
DNA from the genome of a transfected organism, wherein
the genomic DNA of said transfected organism contains a
fragment having two tandemly oriented FLP recombination
target sites therein, said method comprising contacting
25 genomic DNA from said organism with FLP.

25. A method for the precisely targeted
integration of DNA into the genome of a host organism,
said method comprising:
30 (i) introducing a FLP recombination target
site into the genome of cells which are
compatible with the cells of the subject,

-39-

- (ii) introducing a first DNA comprising a nucleotide sequence containing at least one FLP recombination target site therein into the FLP recombination target site in the genome of said cells by contacting said cells with said first DNA and FLP recombinase, and thereafter
- (iii) introducing the cells produced by the process of step (ii) into said subject.

26. A method according to Claim 25, further comprising contacting the genomic DNA from said subject with FLP, thereby recovering the transfected DNA containing said first gene of interest from the genome of said transfected organism.

27. A method according to Claim 26, further comprising introducing at least a portion of a second gene of interest into said FLP recombination target site.

28. A method according to Claim 25, further comprising introducing at least a portion of a second gene of interest into one of the FLP recombination target sites of said subject.

29. A mammalian cell, wherein the genomic DNA of said cell contains at least one FLP recombination target site therein.

30. A mammalian cell according to Claim 29 wherein said FLP recombination target site in the genomic DNA of said cell is positioned within at least a portion of one or more gene(s) of interest.

-40-

31. A mammalian cell according to Claim 30, further comprising DNA encoding, and capable of expressing, in mammalian cells, a FLP recombinase.

5 32. A mammalian cell according to Claim 30 wherein said gene(s) of interest provide a readily analyzable marker feature to the host system.

33. A mammalian cell according to Claim 29
10 wherein said FLP recombination target site has the sequence:

5'-GAAGTTCCTATTCTCTAGAAAGTATAGGAACTT,

15 or functional equivalents thereof.

34. A mammalian cell according to Claim 30 further comprising an additional DNA fragment, wherein said additional DNA fragment is selected from:

- 20 (a) at least a second portion of said first gene of interest, or
 (b) at least a portion of a second gene of interest;

wherein said second DNA contains at least one FLP
25 recombination target site; and wherein said second DNA, when combined in reading frame with said first DNA, provides a functional gene.

35. A transgenic, non-human mammal, wherein
30 said mammal contains at least one FLP recombination target site in the genomic DNA thereof.

-41-

36. A transgenic, non-human mammal according to Claim 35 wherein said FLP recombination target site is positioned within at least a portion of one or more gene(s) of interest.

5

37. A transgenic, non-human mammal according to Claim 35, further comprising a nucleotide sequence encoding, and capable of expressing, in transgenic, non-human mammals, a FLP recombinase.

10

38. A transgenic, non-human mammal according to Claim 35, further comprising FLP recombinase.

39. A transgenic, non-human mammal according to Claim 36 wherein said gene(s) of interest provide a readily analyzable marker feature to the host system.

15

40. A transgenic, non-human mammal according to Claim 35 wherein said FLP recombination target site has the sequence:

20

5'-GAAGTTCCTATTCTCTAGAAAGTATAGGAACTT,

or functional equivalents thereof.

25

41. A transgenic, non-human mammal according to Claim 36 further comprising an additional DNA fragment, wherein said additional DNA fragment is selected from:

- (a) at least a second portion of said first gene of interest, or
- (b) at least a portion of a second gene of interest;

30

wherein said second DNA contains at least one FLP recombination target site; and wherein said second DNA, when combined in reading frame with said first DNA, provides a functional gene.

35

42. A method for the site-specific integration of transfected DNA into the genome of a cell according to Claim 29, said method comprising:

- 5 (i) contacting said genome with:
 (a) FLP recombinase, and
 (b) a first DNA comprising at least a portion of a first gene of interest;
 wherein said first DNA contains at least one FLP recombination target site;
10 and thereafter
 (ii) maintaining the product of step (i) under conditions suitable for site-specific integration of said DNA sequence to occur at the FLP recombination target site in
15 said genome of the host cells.

43. A method according to Claim 42 wherein said FLP recombination target site in the genomic DNA of said cell is positioned within at least a portion of one or
20 more gene(s) of interest.

44. A method according to Claim 42 further comprising additionally contacting said host cell with a second DNA, wherein said second DNA is selected from:
25 (a) at least a second portion of said first gene of interest, or
 (b) at least a portion of a second gene of interest;
wherein said second DNA contains at least one FLP
30 recombination target site; and wherein said second DNA, when combined in reading frame with said first DNA, provides a functional gene.

45. A method according to Claim 42 wherein said
35 FLP recombinase is provided by a FLP expression vector.

-43-

46. A method according to Claim 45 wherein the expression of FLP recombinase by said FLP expression vector is subject to regulatory control.

5 47. A method according to Claim 42 wherein said FLP recombination target site is introduced into the genome of said host mammalian cell by transfecting said host cell with a DNA fragment containing at least one recombination target site therein.

10 48. A method according to Claim 42 wherein the FLP recombination target site in the genomic DNA of said host mammalian cell is so positioned that the introduction of additional DNA sequences therein will
15 inactivate the target gene.

 49. A method for the site-specific integration of transfected DNA into the genome of a host according to Claim 35, said method comprising:

- 20 (i) contacting said genome with:
- (a) FLP recombinase, and
 - (b) a first DNA comprising a nucleotide sequence containing at least one FLP recombination target site therein;
- 25 and thereafter
- (ii) maintaining the product of step (i) under conditions suitable for site-specific integration of said DNA sequence to occur at the FLP recombination target site in
30 said genome of the host.

 50. A method according to Claim 49 wherein said FLP recombination target site is positioned within at least a portion of one or more gene(s) of interest.

35

51. A method according to Claim 49 further comprising additionally contacting said host with a second DNA, wherein said second DNA is selected from:

- 5 (a) at least a second portion of said first gene of interest, or
(b) at least a portion of a second gene of interest;

wherein said second DNA contains at least one FLP recombination target site; and wherein said second DNA,
10 when combined in reading frame with said first DNA, provides a functional gene.

52. A method according to Claim 49 wherein said FLP recombinase is provided by a FLP expression vector.

15

53. A method according to Claim 52 wherein the expression of FLP recombinase by said FLP expression vector is subject to regulatory control.

20

54. A method according to Claim 49 wherein said FLP recombination target site is introduced into the genome of said host mammal by transfecting said host with a DNA fragment containing at least one recombination target site therein.

25

55. A method according to Claim 49 wherein the DNA of said host mammal contains at least one FLP recombination target site, and wherein said FLP recombination target site is so positioned that the
30 introduction of additional DNA sequences therein will inactivate the target gene.

56. A method for the analysis of the development of a mammal, said method comprising:

(a) providing a transgenic mammal comprising:

- 5 (i) an expression construct encoding FLP under the control of a conditional promoter, and
- 10 (ii) a reporter construct under the control of the same or a different promoter, wherein said reporter construct encodes a functional or non-functional reporter gene product, and wherein said construct contains at least one FLP recombination target site therein,
- 15 wherein the functional expression of the functional reporter gene is disrupted when said FLP recombination event occurs, or
- 20 wherein the functional expression of the non-functional reporter gene commences when said FLP recombination event occurs; and

(b) following the development of said mammal to determine when expression of functional reporter gene product either commences or is disrupted.

25

57. A method according to Claim 56 wherein said conditional promoter is developmentally-regulated.

30 58. A co-transfection assay for the occurrence of FLP-mediated recombination, said assay comprising:

(a) co-transfecting a host mammalian cell with:

- 35 (i) a FLP expression plasmid, and
- (ii) a reporter plasmid comprising a non-functional reporter gene wherein said non-functional reporter gene is inactivated by

-46-

the presence of extraneous DNA containing
at least one recombination target site;
and

5 (b) monitoring said host cell under a variety
of conditions for the gain of expression of functional
reporter gene product.

59. A co-transfection assay for the occurrence
of FLP-mediated recombination, said assay comprising:

10 (a) co-transfecting a host mammalian cell
with:

(i) a FLP expression plasmid, and
(ii) a reporter plasmid comprising a
functional reporter gene containing at
15 least one recombination target site
therein; and

(b) monitoring said host cell under a variety
of conditions for the loss of expression of functional
reporter gene product.

20

FIG. 1A







FIG. 1B




SUBSTITUTE SHEET

FIG.2A

pFRT β GAL




pNEO β GAL



0 0.1 0.2 0.5 1 2 6 12 18
MICROGRAMS pOG44

FIG.2C



pOG44 FLP






FIG.3A

E25



E25B



B2N





FIG.3B

E25 E25 E25 B2 B2
B1 B2 N1 N2



INTERNATIONAL SEARCH REPORT

International Application No. PCT/US92/01899

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ³ According to International Patent Classification (IPC) or to both National Classification and IPC IPC (5): C12N 15/85, 5/16; C07H 15/12 US CL : 435/320.1, 240.2, 172.3; 536/27						
II. FIELDS SEARCHED <div style="text-align: center; border: 1px solid black; padding: 2px; margin: 5px 0;">Minimum Documentation Searched⁴</div> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 20%; padding: 5px;">Classification System</th> <th style="padding: 5px;">Classification Symbols</th> </tr> <tr> <td style="padding: 5px; text-align: center;">U.S.</td> <td style="padding: 5px;">435/320.1, 240.2, 172.3; 536/27</td> </tr> </table> <div style="text-align: center; border: 1px solid black; padding: 2px; margin: 5px 0;">Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched⁵</div> <p style="padding: 5px;">Please See Attached Sheet.</p>			Classification System	Classification Symbols	U.S.	435/320.1, 240.2, 172.3; 536/27
Classification System	Classification Symbols					
U.S.	435/320.1, 240.2, 172.3; 536/27					
III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴						
Category*	Citation of Document, ¹⁶ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸				
Y	Cell, volume 59, issued 03 November 1989, Golic <u>et al.</u> , "The FLP Recombinase Of Yeast Catalyzes Site-Specific Recombination In The Drosophila Genome", pages 499-509. See entire document.	1-18, 29-34				
Y	Cell, volume 21, issued September 1980, Broach <u>et al.</u> , "Replication And Recombination Functions Associated With The Yeast Plasmid, 2 μ Circle", pages 501-508. See entire document.	1-18, 29-34				
Y	Proc. Natl. Acad. Sci. USA, volume 82, issued September 1985, Jayaram, "Two-Micrometer Circle Site-Specific Recombination: The Minimal Substrate And The Possible Role Of Flanking Sequences", pages 5875-5879. See entire document.	1-18, 29-34				
Y	J. Bacteriol., volume 169, no. 12, issued December 1987, Utatsu <u>et al.</u> , "Yeast Plasmids Resembling 2 μ m DNA: Regional Similarities And Diversities At The Molecular Level", pages 5537-5545. See entire document.	1-18, 29-34				
<div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>* Special categories of cited documents:¹⁶</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p> </div> </div>						
IV. CERTIFICATION						
Date of the Actual Completion of the International Search ² <div style="text-align: center; font-weight: bold; font-size: 1.2em;">06 JUNE 1992</div>	Date of Mailing of this International Search Report ² <div style="text-align: center; font-weight: bold; font-size: 1.2em;">24 JUN 1992</div>					
International Searching Authority ¹ <div style="text-align: center; font-weight: bold;">ISA/US</div>	Signature of Authorized Officer ²⁰ <div style="text-align: center;"> <div style="text-align: center; font-weight: bold;">CHRISTOPHER S. F. LOW</div> </div>					

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

Y	Molec. Cell. Biol., volume 7, no. 6, issued June 1987, Sauer, "Functional Expression Of The <u>cre-lox</u> Site-Specific Recombination System In The Yeast <u>Saccharomyces cerevisiae</u> , pages 2087-2096. See entire document.	1-18, 29-34
Y	J. Molec. Biol., volume 201, issued 1988, Senecoff <u>et al.</u> , "DNA Recognition By The FLP Recombinase Of The Yeast 2 μ Plasmid. A Mutational Analysis Of The FLP Binding Site", pages 405-421. See entire document.	1-18, 29-34
Y	US, A, 4,959,317 (Sauer) issued 25 September 1990. See at least col 14.	1-18, 29-34

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE¹

1. ☐ Claim numbers , because they relate to subject matter (1) not required to be searched by this Authority, namely:
2. ☐ Claim numbers , because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out (1), specifically:
3. ☐ Claim numbers , because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING²

This International Searching Authority found multiple inventions in this international application as follows:
Please See Attached Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
1-18 and 29-34
(Telephone Practice)
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Search Authority did not invite payment of any additional fee.

Remark on protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PREVIOUS SHEETS

II. FIELDS SEARCHED

Other Documents Searched:

APS: USPAT and JPOABS

DIALOG: BIOSIS, MEDLINE, CA SEARCH, SCISEARCH, LIFE SCIENCES COLLECTION, PASCAL, CONFERENCE PAPERS INDEX, WORLD PATENTS INDEX

SEARCH TERMS: FLP recombina?

VI. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked so as to form single inventive concept and does not meet the criteria of PCT Rule 13.1 and 13.2. The sequential Roman numerals identify the groups.

- I. Claims 1-18 and 29-34, drawn to a mammalian recombination system, DNA, transformed host cells, and method of assembling functional genes are classified in Class 435, subclasses 320.1, 240.2, 172.3 and Class 536, subclass 27.
- II. Claims 19-23, drawn to a method of disrupting gene function are classified in Class 435, subclass 172.3.
- III. Claim 24, drawn to a method of recovering DNA is classified in Class 435, subclass 91.
- IV. Claims 25-28 and 42-55, drawn to methods of targeted (site specific) DNA integration contain species drawn to cells per se (claims 25-28 and 42-48) and claims drawn to non-human transgenic animals (claims 25-28 and 49-55) are classified in Class 435, subclasses 172.1, 172.3, 240.2 and in Class 800 subclass 2.
- V. Claims 35-41, drawn to non-human transgenic animals are classified in Class 800, subclass 2.
- VI. Claims 56 and 57, drawn to a method for analysis of mammalian development are classified in Class 435, subclass 4.
- VII. Claims 58 and 59, drawn to a cotransfection assay are classified in Class 435, subclass 6.

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category*	Citation of Document, ¹⁶ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
Y	Proc. Natl. Acad. Sci. USA, volume 85, issued July 1988, Sauer <u>et al.</u> , "Site-Specific DNA Recombination In Mammalian Cells By The Cre Recombinase Of Bacteriophage P1", pages 5166-5170.	1-18, 29-34
Y	Nucleic Acids Research, volume 17, no. 1, issued 11 January 1989, Sauer <u>et al.</u> , "Cre-Stimulated Recombination At <u>loxP</u> -Containing DNA Sequences Placed Into The Mammalian Genome", pages 147-161. See entire document.	1-18, 29-34
Y	The New Biologist, volume 2, no. 5, issued May 1990, Sauer <u>et al.</u> , "Targeted Insertion Of Exogenous DNA Into The Eukaryotic Genome By The Cre Recombinase", pages 441-449. See entire document.	1-18, 29-34
Y	US, A, 4,997,757 (Schiestl) issued 05 March 1991. See entire document.	1-18, 29-34
A,X	Science, volume 251, issued 15 March 1991, O'Gorman <u>et al.</u> , "Recombinase-Mediated Gene Activation and Site-Specific Integration In Mammalian Cells", pages 1351-1355. See entire document.	1-18, 29-34




FIG. 1A





FIG. 1B




SUBSTITUTE SHEET


2 / 3

FIG.2A

pFRTB_{GAL}




pNEOB_{GAL}



pOG44 FLP



FIG.2B



0 0.1 0.2 0.5 1 2 6 12 18
MICROGRAMS pOG44

FIG.2C




FIG.3A




FIG.3B

